

CERN-PH-EP-2012-337
February 19, 2013

Charged kaon femtoscopic correlations in pp collisions at $\sqrt{s} = 7$ TeV

The ALICE Collaboration*

Abstract

Correlations of two charged identical kaons ($K^{\text{ch}}K^{\text{ch}}$) are measured in pp collisions at $\sqrt{s} = 7$ TeV by the ALICE experiment at the Large Hadron Collider (LHC). One-dimensional $K^{\text{ch}}K^{\text{ch}}$ correlation functions are constructed in three multiplicity and four transverse momentum ranges. The $K^{\text{ch}}K^{\text{ch}}$ femtoscopic source parameters R and λ are extracted. The $K^{\text{ch}}K^{\text{ch}}$ correlations show a slight increase of femtoscopic radii with increasing multiplicity and a slight decrease of radii with increasing transverse momentum. These trends are similar to the ones observed for $\pi\pi$ and $K_s^0K_s^0$ correlations in pp and heavy-ion collisions. However at high multiplicities, there is an indication that the one-dimensional correlation radii for charged kaons are larger than those for pions in contrast to what was observed in heavy-ion collisions at RHIC.

*See Appendix A for the list of collaboration members

1 Introduction.

Extremely high energy densities achieved in heavy-ion collisions at the Large Hadron Collider (LHC) may entail the formation of the Quark-Gluon Plasma (QGP), a state characterized by partonic degrees of freedom [1]. Studying the QGP is the main goal of the ALICE experiment (A Large Ion Collider Experiment) [2]. The system created in ultrarelativistic pp collisions at LHC energies might be similar to the system created in non-central heavy-ion collisions because of the large energy deposited in the overlapping region and therefore may also manifest a collective behavior. The highly compressed strongly-interacting system is expected to undergo longitudinal and transverse expansions. Experimentally, the expansion and the spatial extent at decoupling are observable via Bose-Einstein correlations.

Bose-Einstein correlations of two identical pions at low relative momenta were first shown to be sensitive to the spatial scale of the emitting source by G. Goldhaber, S. Goldhaber, W. Lee and A. Pais 50 years ago [3]. The correlation method since developed and known at present as “correlation femtoscopy” was successfully applied to the measurement of the space-time characteristics of particle production processes in high energy collisions, especially in heavy-ion collisions (see, e.g. [4, 5, 6]). Bose-Einstein correlations of identical particles were widely studied in heavy-ion collisions at the Relativistic Heavy-Ion Collider (RHIC) [7], and were found to confirm the hydrodynamic type of collective expansion of the fireball created in such collisions. In heavy-ion collisions the decrease of the correlation radii with increasing particle momentum was usually considered as a manifestation of a collective behavior of the matter created in such collisions [6]. Event multiplicities reached in 7 TeV pp collisions at the LHC are comparable with those measured in peripheral A+A collisions at RHIC, making the study of the particle momentum dependence of the correlation radii an important test of the collectivity in pp collisions.

The ALICE Collaboration has already studied two-pion correlation radii in pp collisions at 900 GeV [8] and 7 TeV [9], and $K_s^0 K_s^0$ correlation radii in pp collisions at 7 TeV [10]. Two-pion Bose-Einstein correlations in pp collisions at $\sqrt{s} = 900$ GeV and 7 TeV have been successfully described within the EPOS+hydro model [11]. It was shown that the hydrodynamic expansion substantially modifies the source evolution compared to the “classical” EPOS scenario with independent decay of flux-tube strings, allowing one to describe the transverse momentum dependence of the correlation radii at high multiplicities.

The main motivations for carrying out the present $K^{\text{ch}} K^{\text{ch}} (K^+ K^+ + K^- K^-)$ femtoscopy analysis are: 1) study the transverse mass, m_T , dependence of the correlation radii (“ m_T -scaling” is expected to be an additional confirmation of the hydrodynamic type of expansion [6]), 2) get a clearer signal (kaons are less affected by the decay of resonances than pions).

Previous $K^{\text{ch}} K^{\text{ch}}$ studies carried out in Pb–Pb collisions at SPS by the NA44 and NA49 Collaborations [12] and in Au–Au collisions at RHIC by the PHENIX Collaboration [13] revealed scaling in transverse mass: the source sizes versus m_T for different particle types (π , K) fall on the same curve.

$K^{\text{ch}} K^{\text{ch}}$ studies were performed in a combined data from $\alpha\alpha$, pp, and $p\bar{p}$ collisions at ISR by AFS Collaboration [14], in e^+e^- collisions at LEP by the OPAL and DELPHI Collaborations [15], in ep collisions by the ZEUS Collaboration [16]. Due to statistics limitations, only one-dimensional radii were extracted in these experiments, no multiplicity and transverse momentum studies were performed.

In this article we present the measurements of Bose-Einstein correlations for charged kaons in pp collisions at $\sqrt{s} = 7$ TeV performed by the ALICE Collaboration at the LHC. The present study is the first femtoscopic $K^{\text{ch}} K^{\text{ch}}$ study to be carried out in pp collisions and in more than one multiplicity and pair transverse momentum, k_T , range.

The paper is organized as follows: in Section II we describe the ALICE experimental setup and data taking conditions for the data sample used in this work. In Section III we present the correlation measurements and the correlation functions. In Section IV we show the main results obtained in this work:

the one-dimensional radii extracted from the data. We discuss various observed features and compare the results with $K_s^0 K_s^0$ and $\pi\pi$ radii previously measured by the ALICE Collaboration. Finally in Section V we summarize our results.

2 Data analysis.

Approximately 300 million minimum-bias events at $\sqrt{s} = 7$ TeV, recorded in 2010, were analyzed. The ALICE Time Projection Chamber (TPC) and Inner Tracking System (ITS) were used for charged particle track reconstruction and the determination of the primary vertex of the collision.

The TPC identifies charged particles according to their ionization trajectories in the Ne – CO₂ gas. The ionization electrons drift up to 2.5 m from the central electrode to the end caps to be measured on 159 padrows, grouped into 18 sectors. The position at which the track crosses the padrow is determined with a resolution of 2 mm and 3 mm in the drift and transverse directions, respectively. The ITS consists of six silicon layers, two innermost Silicon Pixel Detector (SPD) layers, two Silicon Drift Detector (SDD) layers, and two outer Silicon Strip Detector (SSD) layers, which provide up to six space points for each track.

The forward scintillator detectors VZERO were included in the minimum-bias trigger and their timing signal was used to reject the beam-gas and beam-halo collisions. The minimum-bias trigger required a hit in one of the VZERO counters or in one of the two inner layers of the SPD. The VZERO detectors are placed along the beam line at +3 m and -0.9 m from the nominal interaction point. They cover a region $2.8 < \eta < 5.1$ and $-3.7 < \eta < -1.7$.

ALICE provides excellent particle identification capabilities, using the measurement of specific particle energy loss (dE/dx) in the TPC and the ITS and the time-of-flight (t_{TOF})

information obtained in the Time-Of-Flight (TOF) detector. The TOF detector is based on Multi-gap Resistive Plate Chambers (MRPCs) in a cylindrical configuration at radius 370-399 cm from the beam axis, with about 153.000 readout channels of dimension 3.5×2.5 cm². The start time of the collision (event time zero) is measured by the T0 detector, an array of Cherenkov counters located at +350 cm and -70 cm along the beam-line. If the T0 signal is absent the start time is estimated from the particle arrival times at the TOF. The overall time-of-flight resolution depends on the TOF timing signal resolution (better than 100 ps), the accuracy of the reconstructed flight path and the uncertainty in the event start time. The resulting time-of-flight resolution is about 160 ps.

The following criteria were applied for the event selection:

- z -position of the reconstructed vertex within ± 10 cm around the geometrical center of the ALICE detector;
- at least one particle in the event reconstructed and identified as a charged kaon (*In fact, the correlation signal is constructed from events having at least two same-charged kaons (a pair). The one-kaon events do contribute to the mixed background. It was verified that including one-kaon event to the mixed background does not change the shape of the correlation function.*)

The criteria for track selection are listed below:

- the kaons were selected in the kinematic ranges: $|\eta| < 1.0$ and $0.15 < p_T < 1.2$ GeV/c;
- tracks must include at least 70 space points (or clusters) out of a maximum possible number 159 in the TPC, and two space points in the ITS (of maximum 6).

- the quality of a track was determined by the χ^2/N value for the Kalman fit to the constructed position of the TPC clusters (N is the number of clusters associated with the track); the track was rejected if the value was larger than 4.0 (2 degrees of freedom per cluster);
- in order to reduce the number of secondary particles it was required that the particle trajectory distance from the primary vertex was less than 0.2 cm in the transverse plane and less than 0.25 cm in the beam direction.

Usually the femtoscopic correlation functions of identical particles are very sensitive to two-track reconstruction effects because particles have close momenta and thus close trajectories. The “splitting” of the tracks means that one track was reconstructed as two, and “merging” means that two different tracks were reconstructed as one. For the correlation structures measured in pp collisions, with characteristic widths ~ 0.2 GeV/ c , track splitting and track merging in the event reconstruction are small effects, but we applied the standard femtoscopic double track cuts (see for details [9]):

- “anti-splitting cut”: pairs which share more than 5% of clusters in the TPC were removed;
- “anti-merging cut”: pairs that are separated by less than 3 cm at the entrance of the TPC were removed.

Pair cuts were applied in exactly the same way for real (signal) and mixed (background) pairs.

In the present analysis the limit $p_T < 1.2$ GeV/ c for kaon selection on the TPC and TOF signals was used in order to ensure a high purity of the kaon sample. Kaons were selected by requiring that the deviation of the specific (dE/dx) energy loss in the TPC from that calculated with a parametrized Bethe-Bloch formula be within some number of sigma standard deviations ($N_{\sigma_{\text{TPC}}}$). A similar $N_{\sigma_{\text{TOF}}}$ method was applied for the particle identification in the TOF using the difference between the measured time-of-flight and the calculated one as a function of the track length and the particle momentum at each tracking step and for each particle mass.

More details on the particle identification are given in Ref. [17], where it is shown in particular that the fraction efficiency of the particles reconstructed by the TPC with associated signal in the TOF (the TOF matching efficiency) deviates from 30% up to 55% in the p_T -region – under study in our paper. In the present analysis strict cuts on TPC and TOF signals for kaon selection were used in order to provide the better purity of the kaon sample. The relative contribution of the kaons from the sample with TOF signal to the full sample of the identified kaons used in this analysis is about 60%.

In the present analysis strict cuts on TPC and TOF signals for kaon selection were used in order to provide the better purity of the kaon sample. If the TOF signal was not available, the following cuts were taken:

- $N_{\sigma_{\text{TPC}}} < 1$ for $p < 0.35$ GeV/ c ;
- $N_{\sigma_{\text{TPC}}} < 2$ for $0.35 < p < 0.6$ GeV/ c ;
- at larger momenta the tracks were rejected because of significant pion contamination.

If the TOF signal was available, we required that $N_{\sigma_{\text{TOF}}} < 3$ and $N_{\sigma_{\text{TPC}}} < 3$.

Figure 1 shows the transverse momentum dependence of the kaon purity (the ratio of correctly identified kaons to all identified ones) obtained with Monte Carlo (PYTHIA) simulations. The contamination comes mainly from e^+/e^- with maximum $\sim 25\%$ for $0.35 < p_T < 0.6$ GeV/ c and also from pions at the level $\sim 1 - 3\%$ for $0.35 < p_T < 1.2$ GeV/ c . In consequence, the probability of selecting an ee or




Fig. 1: Purity and contaminations of selected kaons in pp collisions at $\sqrt{s} = 7$ TeV.

a $\pi\pi$ pair instead of a KK pair, even in the case where the contamination is maximal, is still rather low, smaller than $\sim 6\%$. The probability of selecting an eK(Ke) pair instead of a KK pair can reach $\sim 37\%$ for $0.35 < p_T < 0.6$ GeV/c. The probability of selecting a $\pi K(K\pi)$ pair instead of a KK pair is less than $\sim 6\%$ for $0.35 < p_T < 1.2$ GeV/c. Note that such contamination modifies only the strength of correlation and not the shape of the correlation function. Using the Monte Carlo information and also the purity of kaons estimated from the measured TPC dE/dx distributions and a realistic single-particle kinematics we have estimated the purity of kaon pairs at low relative momentum as: $p = 0.84 \pm 0.05$ (stat.) ± 0.15 (syst.), 0.61 ± 0.03 (stat.) ± 0.12 (syst.), 0.79 ± 0.04 (stat.) ± 0.07 (syst.), 1.0 ± 0.05 (stat.) ± 0.05 (syst.) for pair transverse momentum $k_T = |\mathbf{p}_{T,1} + \mathbf{p}_{T,2}|/2$: (0.2-0.35), (0.35-0.5), (0.5-0.7), (0.7-1.0) GeV/c, respectively.

3 Charged kaon correlation functions.

Momentum correlations are usually studied by means of correlation functions of two or more particles. Specifically, the two-particle correlation function $CF(\mathbf{p}_1, \mathbf{p}_2) = A(\mathbf{p}_1, \mathbf{p}_2)/B(\mathbf{p}_1, \mathbf{p}_2)$ is defined as the ratio of the two-particle distribution in a given event $A(\mathbf{p}_1, \mathbf{p}_2)$ to the reference one, $B(\mathbf{p}_1, \mathbf{p}_2)$, where \mathbf{p}_1 and \mathbf{p}_2 are the momentum vectors of the two particles. In the present analysis the reference distribution is constructed by mixing particles of a given class, as described below.

The analysis was performed in three multiplicity ranges based on the measured charged-particle multiplicity, N_{ch} : (1 – 11), (12 – 22), (> 22), and in 4 ranges of pair transverse momentum k_T : (0.2-0.35), (0.35-0.5), (0.5-0.7), (0.7-1.0) GeV/c. Event multiplicity was determined as the number of charged particles emitted into the pseudorapidity range $|\eta| < 1$ and transverse momentum range $0.12 < p_T < 10$ GeV/c. For each class of events we calculated the charged-particle pseudorapidity density $dN_{ch}/d\eta$ corrected for the detection efficiency obtained with Monte Carlo. The considered event multiplicity ranges (1 – 11), (12 – 22), > 22 , correspond to mean charged particle densities, $dN_{ch}/d\eta$, of 3.2, 8.1 and 17.2, respectively, with systematic uncertainties of $\sim 5\%$.

The numerators and denominators of positive and negative kaon distributions were summed up before constructing the ratio (K^+K^+ and K^-K^- correlation functions were found to coincide within errors thus justifying the procedure). The function is normalized to unity in the range $0.5 < q_{inv} < 1.0$ GeV/c, where

$q_{\text{inv}} = \sqrt{|\mathbf{q}|^2 - q_0^2}$, $\mathbf{q} = \mathbf{p}_1 - \mathbf{p}_2$, and $q_0 = E_1 - E_2$. The range for normalization was chosen outside the Bose-Einstein peak.

The correlation function is fitted by a single-Gaussian [18]:

$$CF(q_{\text{inv}}) = (1 - \lambda + K(q_{\text{inv}}) (\lambda \exp(-R_{\text{inv}}^2 q_{\text{inv}}^2))) D(q_{\text{inv}}), \quad (1)$$

where the factor $K(q_{\text{inv}})$ is the Coulomb function integrated over a spherical source of 1 fm. The function $D(q_{\text{inv}})$, “baseline”, takes into account all non-femtoscopic correlations, including the long-range correlations due to energy-momentum conservation. The parameters R_{inv} and λ describe the size of the kaon source, and the correlation strength, respectively. The λ -parameter depends also on purity and decreases if the purity is not 100%. The R_{inv} is measured in the pair rest frame.

The baseline was fitted by a standard quadratic polynomial

$$D(q_{\text{inv}}) = 1 + a q_{\text{inv}} + b q_{\text{inv}}^2. \quad (2)$$

To estimate the systematic errors due to the fitting procedure, other functions with derivatives equal to zero at $q_{\text{inv}} = 0$ were also employed, such as:

$$D(q_{\text{inv}}) = \sqrt{1 + a q_{\text{inv}}^2 + b q_{\text{inv}}^4} \quad (3)$$

and the Gaussian:

$$D(q_{\text{inv}}) = b(1 + \exp(-a q_{\text{inv}}^2)). \quad (4)$$

The PERUGIA-2011 tune [19] of the Monte Carlo event generator PYTHIA [20] describes well the kaon spectra in pp collisions at LHC energies. Therefore, it was used to simulate the correlation function without the Bose-Einstein effect.

4 Results and Discussion.

Figure 2 presents the experimental two-kaon correlation functions and those obtained from a simulation using PERUGIA-2011 (open circles) as a function of the invariant pair relative momentum. As one can see, the Monte Carlo simulation reproduces well the experimental correlation function at large q_{inv} , i.e. the long-range correlations. The model does not contain the Bose-Einstein effect, so the enhancement at low q_{inv} is due to non-femtoscopic correlations in PYTHIA, probably arising from mini-jets. The baseline points, obtained from PERUGIA-2011 were fitted to Eq. (2). The parameters a and b were used in the fitting of the experimental points by Eq. (1). The same method was used to model the baseline for the ALICE $\pi\pi$ correlation studies in 0.9 [8] and 7 TeV [9] pp collisions. Figures 3-4 and Table 1 present the one-dimensional λ -parameters and Gaussian radii versus $m_T = \sqrt{k_T^2 + m_K^2}$ including statistical and systematical errors.

In order to estimate the systematic error from the choice of baseline functional form we repeated the fitting procedure using the baseline fitted with Eqs. (3-4). The radii obtained in the three ways differ by less than 4 % at low multiplicities (or k_T) and by up to 10% at high multiplicities (or k_T). The systematic errors estimated from varying the q_{inv} fit range are below 2 % and up to 15% at low and high multiplicities (or k_T) bins respectively. The systematic errors from splitting and merging effects were estimated by using different start points for the fit of correlation function: 0.03 and 0.06 GeV/c, and its are about 2-6%. The systematic error connected to the Coulomb function in Eq. (1) was calculated in the following way: at first, the radius of the spherical source was taken equal to 1 fm, then the fitting procedure was repeated using these radii $\pm 3\sigma_R$ (where σ_R is the total error) as the argument of the Coulomb function. The obtained systematic error is about 2-4%.




Fig. 2: Correlation functions versus q_{inv} for identical kaons from pp collisions at $\sqrt{s} = 7$ TeV (solid circles) and those obtained with PERUGIA-2011 (open circles). Positive and negative kaon pairs are combined. The three columns represent the samples with different charged-particle multiplicities: (1 – 11), (12 – 22), (> 22), the four rows represent the four pair transverse momentum ranges: (0.2-0.35), (0.35-0.5), (0.5-0.7), (0.7-1.0) GeV/c. The lines going through the points represent the Gaussian fits discussed in the text.




Fig. 3: λ -parameters of $K^{\text{ch}}K^{\text{ch}}$ versus m_T extracted by fitting correlation functions shown in Fig. 2 to Eq. (1) and the baseline to Eq. (2). For comparison the $K_s^0K_s^0$ [10] λ -parameters measured by ALICE in 7 TeV pp collisions are also shown. Statistical (darker lines) and total errors are shown. The points corresponding to the second and third multiplicity bins are offset by $0.03 \text{ GeV}/c^2$ for clarity.

The m_T dependence of λ shown in Fig. 3 demonstrates that λ varies within the range $\sim 0.3 - 0.5$ (except the first point at lowest multiplicity and lowest k_T , which is ~ 0.2). As seen in Fig. 3, the λ -parameters for $K^{\text{ch}}K^{\text{ch}}$ are generally smaller than those for $K_s^0K_s^0$ [10]. There are several reason for the λ -parameter to be less than the ideal case of unity. Possible causes may be such as a partially coherent source, a contribution to the observed kaons from decays of long-lived resonances, a deviation from the Gaussian parameterization due to a mixture of sources with different radii (see e.g. [21]), or a particle misidentification. The latter cause influences mostly the $K^{\text{ch}}K^{\text{ch}}$ sample in the momentum range $0.4 < p_T < 0.6 \text{ GeV}/c$ (Fig. 1). For the future, it seems desirable to improve the PID and thus increase the purity and to improve the description of non-femtoscopic correlations using different models.

The $K^{\text{ch}}K^{\text{ch}}$ correlation radii in Fig. 4 show an increase with multiplicity in agreement with the $\pi\pi$ radii at 900 GeV [8], and 7 TeV [9], and the $K_s^0K_s^0$ radii [10], as it was observed for $\pi\pi$ correlations in heavy-ion collisions [22]. These radii also decrease with increasing m_T for the large multiplicity bins N_{ch} (12 – 22) and $N_{\text{ch}} (> 22)$. Such a tendency was found for pions [9] and neutral kaons $K_s^0K_s^0$ [10] in pp collisions, and pions in heavy-ion collisions at LHC energies [22]. In the low multiplicity bin N_{ch} (1 – 11) charged kaons show a completely different k_T -dependence of the radii: these radii increase with k_T . This effect is qualitatively similar to that of pions [9].

It was observed that λ -parameters (Fig. 3) are correlated with the radii (Fig. 4). Such a correlation can result from the non-perfect fit results if the fit quality depends on m_T . The reasons of such a dependence are: non ideal description of the baseline by PERUGIA-2011, especially at large k_T , non-Gaussian shape of the source due to resonance contribution and non-spherical shape of the source. The last two points mean that our one-dimensional Gaussian fit is only an approximate description of the source. In pp collisions the effect of this non-Gaussian shape of the correlation function due to different sizes in the 'x-y-z' directions plays a more important role than in heavy-ion collisions. This requires a detailed 3D analysis, which is foreseen for $K^{\text{ch}}K^{\text{ch}}$ correlation functions with the new large set of data recorded by the ALICE Collaboration in 2011 and 2012.




Fig. 4: One-dimensional charged kaon radii versus m_T extracted by fitting correlation functions shown in Fig. 2 to Eq. (1) and the baseline to Eq. (2). For comparison the $\pi\pi$ [9] and $K_s^0 K_s^0$ [10] radii measured by ALICE in 7 TeV pp collisions are also shown. Statistical (darker lines) and total errors are shown. The points corresponding to the second and third multiplicity bins are offset by $0.03 \text{ GeV}/c^2$ for clarity.

The m_T dependence of the radii in heavy-ion collisions was interpreted as the manifestation of the strong collective hydrodynamic expansion of the created matter [6]. The observed similar behavior in pp collisions, shown in Fig.4, has some specific features: 1) at low multiplicity the radii increase with k_T , 2) there is no distinct m_T scaling: the kaon radii seem to be larger than the pion ones. The model calculations performed in [11] can successfully describe the different behavior of pion correlation radii in low and high multiplicity bins, suggesting that the contribution of the hydrodynamic phase is negligible in low-multiplicity events, while for events with high multiplicity, it is substantial.

As shown in [23], due to the small size of the created system in pp collisions, the flow of resonances may play a significant role in large multiplicity bins, where essential hydrodynamic collective flow is expected [11]. According to simple chemical model calculations [10], the influence of this flow should be relatively smaller for kaons than for pions, leading to the effect that the kaon radii can be larger than the pion ones. The measured $K^{\text{ch}} K^{\text{ch}}$ correlation radii displayed in Fig.4 support such an hypothesis, however a detailed theoretical study is needed.

5 Summary.

The ALICE Collaboration has measured charged kaon correlation functions in pp collisions at $\sqrt{s} = 7 \text{ TeV}$ at the LHC. In agreement with the previous measurements in pp and heavy-ion collisions at lower energies, the extracted correlation radii R_{inv} increase with the event multiplicity and decrease with the pair transverse mass/momentum. The novel features are some hints to the increase of the radii with m_T in the low-multiplicity bin and to the fact that kaon radii are larger than the pion ones. These peculiarities deserve further experimental and theoretical studies.

Table 1: $K^{\text{ch}}K^{\text{ch}}$ source parameters vs. k_T for $\sqrt{s}=7$ TeV pp collisions. Statistical and systematic errors are listed.

k_T range (GeV/c)	N_{ch}	$dN_{\text{ch}}/d\eta$	$\langle k_T \rangle$ (GeV/c)	λ	R_{inv} (fm)
0.20-0.35	1-11	3.2	0.28 ± 0.04	$0.20 \pm 0.04 \pm 0.03$	$0.47 \pm 0.07 \pm 0.12$
0.35-0.50	1-11	3.2	0.42 ± 0.05	$0.31 \pm 0.03 \pm 0.02$	$0.65 \pm 0.05 \pm 0.06$
0.50-0.70	1-11	3.2	0.59 ± 0.06	$0.39 \pm 0.08 \pm 0.03$	$0.91 \pm 0.10 \pm 0.06$
0.07-1.00	1-11	3.2	0.80 ± 0.08	$0.23 \pm 0.10 \pm 0.20$	$0.81 \pm 0.21 \pm 0.24$
0.20-0.35	12-22	8.1	0.28 ± 0.04	$0.51 \pm 0.12 \pm 0.03$	$1.45 \pm 0.15 \pm 0.02$
0.35-0.50	12-22	8.1	0.42 ± 0.05	$0.46 \pm 0.04 \pm 0.04$	$1.18 \pm 0.06 \pm 0.03$
0.50-0.70	12-22	8.1	0.59 ± 0.06	$0.34 \pm 0.07 \pm 0.10$	$1.05 \pm 0.12 \pm 0.14$
0.70-1.00	12-22	8.1	0.80 ± 0.08	$0.21 \pm 0.04 \pm 0.10$	$0.73 \pm 0.07 \pm 0.10$
0.20-0.35	>22	17.2	0.28 ± 0.04	$0.51 \pm 0.08 \pm 0.03$	$1.53 \pm 0.10 \pm 0.02$
0.35-0.50	>22	17.2	0.42 ± 0.05	$0.44 \pm 0.03 \pm 0.04$	$1.44 \pm 0.04 \pm 0.03$
0.50-0.70	>22	17.2	0.59 ± 0.06	$0.30 \pm 0.03 \pm 0.04$	$1.25 \pm 0.06 \pm 0.06$
0.70-1.00	>22	17.2	0.80 ± 0.08	$0.37 \pm 0.05 \pm 0.06$	$1.31 \pm 0.08 \pm 0.08$

References

- [1] N. Cabibbo, G. Parisi, Phys. Lett. **B59**, 67 (1975) 67; E. V. Shuryak, Phys. Rept. **61** (1980) 71;
- [2] K. Aamodt *et al.* [ALICE Collaboration], JINST **3**, S08002 (2008).
- [3] G. Goldhaber, S. Goldhaber, W.-Y. Lee, and A. Pais, Phys. Rev. **120**, 300 (1960).
- [4] M.I. Podgoretsky, Fiz. Elem. Chast. Atom. Yad. **20**, 628 (1989) [Sov. J. Part. Nucl. **20**, 266 (1989)].
- [5] R. Lednicky, Phys. of Atomic Nuclei **67**, 71 (2004). [arXiv:0903.1296 [nucl-ex]].
- [6] S. Pratt, Phys. Rep. Lett. **53** 1219 (1984); M. Lisa, S. Pratt, R. Soltz, U. Wiedemann, Ann. Rev. Nucl. Part. Sci. **55**, 357 (2005).
- [7] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **93**, 152302 (2004), [nucl-ex/0401003]; B. I. Abelev *et al.* [STAR Collaboration], Phys. Rev. C **80**, 024905 (2009),
- [8] K. Aamodt *et al.* [ALICE Collaboration], Phys. Rev. D **82** 052001 (2010) [arXiv:1007.0516 [hep-th]].
- [9] K. Aamodt *et al.* [ALICE Collaboration], Phys. Rev. D **84**, 112004 (2011) [arXiv:1101.3665 [hep-ex]].
- [10] T. J. Humanic, J. Phys. G **38**, 124058 (2011) [arXiv:1107.0084 [hep-ex]]; B. Abelev *et al.* [ALICE Collaboration], Phys. Lett. B **717**, 151 (2012) [arXiv:1206.2056].
- [11] K. Werner, I. Karpenko, T. Pierog, M. Bleicher and K. Mikhailov, Phys. Rev. C **83**, 044915 (2011) [arXiv:1010.0400 [nucl-th]]; K. Werner, K. Mikhailov, I. Karpenko, T. Pierog, [arXiv:1104.2405 [hep-ph]]
- [12] I.G. Bearden *et al.* [NA44 Collaboration], Phys. Rev. Lett. **87** 112301 (2001); S.V. Afanasiev *et al.* [NA49 Collaboration], Phys. Lett. B **557** 157 (2003).
- [13] S. Afanasiev *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **103**, 142301 (2009) [arXiv:0903.4863 [nucl-ex]].
- [14] T. Akesson *et al.* [The Axial Field Spectrometer Collaboration], Phys. Lett. B **155**, 128 (1985).
- [15] G. Abbiendi *et al.* [OPAL Collaboration], Eur. J. Phys. C **21**, 23 (2001), P. Abreu *et al.* [DELPHI Collaboration], Phys. Lett. B **379**, 330 (1996).
- [16] S. Chekanov *et al.* [ZEUS Collaboration], Phys. Lett. B **652**, 1 (2007).
- [17] K. Aamodt *et al.* [ALICE Collaboration], Eur. Phys. J. C **71**, 1655 (2011) [arXiv:1101.4110 [hep-ex]].
- [18] M. G. Bowler, Phys. Lett. B **270**, 69 (1991), Y. Sinyukov, R. Lednicky, S. V. Akkelin, J. Pluta, B. Erasmus, Phys. Lett. B **432**, 248 (1998).

- [19] P. Z. Skands, Phys. Rev. D **82**, 074018 (2010)
- [20] T. Sjostrand, S. Mrenna, and P. Z. Skands, JHEP **05**, 026 (2006), [arXiv:hep-ph/0603175].
- [21] S.V. Afanasiev *et al.* [NA49 Collaboration] Phys. Lett. B **557**, 157 (2003), R. Lednicky and M. I. Podgoretsky, Yad. Fiz. **30** 837 (1979), (Sov. J. Nucl. Phys. **30** 432 (1979)).
- [22] K. Aamodt *et al.* [ALICE Collaboration], Phys. Lett. B **696**, 328 (2011) [arXiv:1012.4035 [nucl-ex]].
- [23] A. Kisiel, Phys. Rev. C **84**, 044913 (2011) [arXiv:1012.1517 [nucl-th]].

6 Acknowledgements

The ALICE collaboration would like to thank all its engineers and technicians for their invaluable contributions to the construction of the experiment and the CERN accelerator teams for the outstanding performance of the LHC complex.

The ALICE collaboration acknowledges the following funding agencies for their support in building and running the ALICE detector:

State Committee of Science, Calouste Gulbenkian Foundation from Lisbon and Swiss Fonds Kidagan, Armenia;

Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Financiadora de Estudos e Projetos (FINEP), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP);

National Natural Science Foundation of China (NSFC), the Chinese Ministry of Education (CMOE) and the Ministry of Science and Technology of China (MSTC);

Ministry of Education and Youth of the Czech Republic;

Danish Natural Science Research Council, the Carlsberg Foundation and the Danish National Research Foundation;

The European Research Council under the European Community's Seventh Framework Programme;

Helsinki Institute of Physics and the Academy of Finland;

French CNRS-IN2P3, the 'Region Pays de Loire', 'Region Alsace', 'Region Auvergne' and CEA, France;

German BMBF and the Helmholtz Association;

General Secretariat for Research and Technology, Ministry of Development, Greece;

Hungarian OTKA and National Office for Research and Technology (NKTH);

Department of Atomic Energy and Department of Science and Technology of the Government of India;

Istituto Nazionale di Fisica Nucleare (INFN) and Centro Fermi - Museo Storico della Fisica e Centro Studi e Ricerche "Enrico Fermi", Italy;

MEXT Grant-in-Aid for Specially Promoted Research, Japan;

Joint Institute for Nuclear Research, Dubna;

National Research Foundation of Korea (NRF);

CONACYT, DGAPA, México, ALFA-EC and the HELEN Program (High-Energy physics Latin-American-European Network);

Stichting voor Fundamenteel Onderzoek der Materie (FOM) and the Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), Netherlands;

Research Council of Norway (NFR);

Polish Ministry of Science and Higher Education;

National Authority for Scientific Research - NASR (Autoritatea Națională pentru Cercetare Științifică - ANCS);

Ministry of Education and Science of Russian Federation, International Science and Technology Center, Russian Academy of Sciences, Russian Federal Agency of Atomic Energy, Russian Federal Agency for Science and Innovations and CERN-INTAS;

Ministry of Education of Slovakia;

Department of Science and Technology, South Africa;
CIEMAT, EELA, Ministerio de Educación y Ciencia of Spain, Xunta de Galicia (Consellería de Educación), CEADEN, Cubaenergía, Cuba, and IAEA (International Atomic Energy Agency);
Swedish Research Council (VR) and Knut & Alice Wallenberg Foundation (KAW);
Ukraine Ministry of Education and Science;
United Kingdom Science and Technology Facilities Council (STFC);
The United States Department of Energy, the United States National Science Foundation, the State of Texas, and the State of Ohio.

A The ALICE Collaboration

B. Abelev⁷², J. Adam³⁷, D. Adamová⁷⁹, A.M. Adare¹²⁸, M.M. Aggarwal⁸³, G. Aglieri Rinella³³, M. Agnello^{99,89}, A.G. Agocs¹²⁷, A. Agostinelli²⁷, Z. Ahammed¹²³, N. Ahmad¹⁷, A. Ahmad Masoodi¹⁷, S.U. Ahn^{40,65}, S.A. Ahn⁶⁵, M. Ajaz¹⁵, A. Akindinov⁵⁰, D. Aleksandrov⁹⁵, B. Alessandro⁹⁹, A. Alici^{101,12}, A. Alkin³, E. Almaráz Aviña⁶⁰, J. Alme³⁵, T. Alt³⁹, V. Altini³¹, S. Altinpinar¹⁸, I. Altsybeev¹²⁴, C. Andrei⁷⁵, A. Andronic⁹², V. Anguelov⁸⁸, J. Anielski⁵⁸, C. Anson¹⁹, T. Antičić⁹³, F. Antinori¹⁰⁴, P. Antonioli¹⁰¹, L. Aphecetche¹⁰⁸, H. Appelshäuser⁵⁶, N. Arbor⁶⁸, S. Arcelli²⁷, A. Arend⁵⁶, N. Armesto¹⁶, R. Arnaldi⁹⁹, T. Aronsson¹²⁸, I.C. Arsene⁹², M. Arslandok⁵⁶, A. Asryan¹²⁴, A. Augustinus³³, R. Averbeck⁹², T.C. Awes⁸⁰, J. Äystö⁴², M.D. Azmi^{17,85}, M. Bach³⁹, A. Badalà¹⁰³, Y.W. Baek^{67,40}, R. Bailhache⁵⁶, R. Bala^{86,99}, R. Baldini Ferroli¹², A. Baldisseri¹⁴, F. Baltasar Dos Santos Pedrosa³³, J. Bán⁵¹, R.C. Baral⁵², R. Barbera²⁶, F. Barile³¹, G.G. Barnaföldi¹²⁷, L.S. Barnby⁹⁷, V. Barret⁶⁷, J. Bartke¹¹¹, M. Basile²⁷, N. Bastid⁶⁷, S. Basu¹²³, B. Bathen⁵⁸, G. Batigne¹⁰⁸, B. Batyunya⁶³, C. Baumann⁵⁶, I.G. Bearden⁷⁷, H. Beck⁵⁶, N.K. Behera⁴⁴, I. Belikov⁶², F. Bellini²⁷, R. Bellwied¹¹⁷, E. Belmont-Moreno⁶⁰, G. Bencedi¹²⁷, S. Beole²⁴, I. Berceanu⁷⁵, A. Bercuci⁷⁵, Y. Berdnikov⁸¹, D. Berenyi¹²⁷, A.A.E. Bergognon¹⁰⁸, D. Berzano^{24,99}, L. Betev³³, A. Bhasin⁸⁶, A.K. Bhati⁸³, J. Bhom¹²¹, L. Bianchi²⁴, N. Bianchi⁶⁹, J. Bielčák³⁷, J. Bielčíková⁷⁹, A. Bilandzic⁷⁷, S. Bjelogrić⁴⁹, F. Blanco¹¹⁷, F. Blanco¹⁰, D. Blau⁹⁵, C. Blume⁵⁶, M. Boccioli³³, S. Böttger⁵⁵, A. Bogdanov⁷³, H. Bøggild⁷⁷, M. Bogolyubsky⁴⁷, L. Boldizsár¹²⁷, M. Bombara³⁸, J. Book⁵⁶, H. Borel¹⁴, A. Borissov¹²⁶, F. Bossú⁸⁵, M. Botje⁷⁸, E. Botta²⁴, E. Braidot⁷¹, P. Braun-Munzinger⁹², M. Bregant¹⁰⁸, T. Breitner⁵⁵, T.A. Browning⁹⁰, M. Broz³⁶, R. Brun³³, E. Bruna^{24,99}, G.E. Bruno³¹, D. Budnikov⁹⁴, H. Buesching⁵⁶, S. Bufalino^{24,99}, P. Buncic³³, O. Busch⁸⁸, Z. Buthelezi⁸⁵, D. Caffarri^{28,104}, X. Cai⁷, H. Caines¹²⁸, E. Calvo Villar⁹⁸, P. Camerini²², V. Canoa Roman¹¹, G. Cara Romeo¹⁰¹, F. Carena³³, W. Carena³³, N. Carlin Filho¹¹⁴, F. Carminati³³, A. Casanova Díaz⁶⁹, J. Castillo Castellanos¹⁴, J.F. Castillo Hernandez⁹², E.A.R. Casula²³, V. Catanescu⁷⁵, C. Cavicchioli³³, C. Ceballos Sanchez⁹, J. Cepila³⁷, P. Cerello⁹⁹, B. Chang^{42,130}, S. Chapeland³³, J.L. Charvet¹⁴, S. Chattopadhyay¹²³, S. Chattopadhyay⁹⁶, I. Chawla⁸³, M. Cherney⁸², C. Cheshkov^{33,116}, B. Cheynis¹¹⁶, V. Chibante Barroso³³, D.D. Chinellato¹¹⁷, P. Chochula³³, M. Chojnacki^{77,49}, S. Choudhury¹²³, P. Christakoglou⁷⁸, C.H. Christensen⁷⁷, P. Christiansen³², T. Chujo¹²¹, S.U. Chung⁹¹, C. Cicalo¹⁰², L. Cifarelli^{27,33,12}, F. Cindolo¹⁰¹, J. Cleymans⁸⁵, F. Coccetti¹², F. Colamaria³¹, D. Colella³¹, A. Collu²³, G. Conesa Balbastre⁶⁸, Z. Conesa del Valle³³, M.E. Connors¹²⁸, G. Contin²², J.G. Contreras¹¹, T.M. Cormier¹²⁶, Y. Corrales Morales²⁴, P. Cortese³⁰, I. Cortés Maldonado², M.R. Cosentino⁷¹, F. Costa³³, M.E. Cotallo¹⁰, E. Crescio¹¹, P. Crochet⁶⁷, E. Cruz Alaniz⁶⁰, E. Cuautle⁵⁹, L. Cunqueiro⁶⁹, A. Dainese^{28,104}, H.H. Dalsgaard⁷⁷, A. Danu⁵⁴, S. Das⁴, I. Das⁴⁶, D. Das⁹⁶, K. Das⁹⁶, A. Dash¹¹⁵, S. Dash⁴⁴, S. De¹²³, G.O.V. de Barros¹¹⁴, A. De Caro^{29,12}, G. de Cataldo¹⁰⁰, J. de Cuveland³⁹, A. De Falco²³, D. De Gruttola²⁹, H. Delagrangé¹⁰⁸, A. Deloff⁷⁴, N. De Marco⁹⁹, E. Dénes¹²⁷, S. De Pasquale²⁹, A. Deppman¹¹⁴, G. D'Erasmus³¹, R. de Rooij⁴⁹, M.A. Diaz Corchero¹⁰, D. Di Bari³¹, T. Dietel⁵⁸, C. Di Giglio³¹, S. Di Liberto¹⁰⁵, A. Di Mauro³³, P. Di Nezza⁶⁹, R. Divià³³, Ø. Djuvsland¹⁸, A. Dobrin^{126,32}, T. Dobrowolski⁷⁴, B. Dönigus⁹², O. Dordic²¹, O. Driga¹⁰⁸, A.K. Dubey¹²³, A. Dubla⁴⁹, L. Ducroux¹¹⁶, P. Dupieux⁶⁷, A.K. Dutta Majumdar⁹⁶, M.R. Dutta Majumdar¹²³, D. Elia¹⁰⁰, D. Emschermann⁵⁸, H. Engel⁵⁵, B. Erasmus^{33,108}, H.A. Erdal³⁵, B. Espagnon⁴⁶, M. Estienne¹⁰⁸, S. Esumi¹²¹, D. Evans⁹⁷, G. Eyyubova²¹, D. Fabris^{28,104}, J. Faivre⁶⁸, D. Falchieri²⁷, A. Fantoni⁶⁹, M. Fasel^{92,88}, R. Fearick⁸⁵, D. Fehlker¹⁸, L. Feldkamp⁵⁸, D. Felea⁵⁴, A. Feliciello⁹⁹, B. Fenton-Olsen⁷¹, G. Feofilov¹²⁴, A. Fernández Téllez², A. Ferretti²⁴, A. Festanti²⁸, J. Figiel¹¹¹, M.A.S. Figueredo¹¹⁴, S. Filchagin⁹⁴, D. Finogeev⁴⁸, F.M. Fionda³¹, E.M. Fiore³¹, E. Floratos⁸⁴, M. Floris³³, S. Foertsch⁸⁵, P. Foka⁹², S. Fokin⁹⁵, E. Fragiaco¹⁰⁶, A. Francescon^{33,28}, U. Frankenfeld⁹², U. Fuchs³³, C. Furget⁶⁸, M. Fusco Girard²⁹, J.J. Gaardhøje⁷⁷, M. Gagliardi²⁴, A. Gago⁹⁸, M. Gallio²⁴, D.R. Gangadharan¹⁹, P. Ganoti⁸⁰, C. Garabatos⁹², E. Garcia-Solis¹³, I. Garishvili⁷², J. Gerhard³⁹, M. Germain¹⁰⁸, C. Geuna¹⁴, M. Gheata^{54,33}, A. Gheata³³, P. Ghosh¹²³, P. Gianotti⁶⁹, M.R. Girard¹²⁵, P. Giubellino³³, E. Gladysz-Dziadus¹¹¹, P. Glässel⁸⁸, R. Gomez^{113,11}, E.G. Ferreira¹⁶, L.H. González-Trueba⁶⁰, P. González-Zamora¹⁰, S. Gorbunov³⁹, A. Goswami⁸⁷, S. Gotovac¹¹⁰, L.K. Graczykowski¹²⁵, R. Grajcarek⁸⁸, A. Grelli⁴⁹, C. Grigoras³³, A. Grigoras³³, V. Grigoriev⁷³, S. Grigoryan⁶³, A. Grigoryan¹, B. Grinyov³, N. Grion¹⁰⁶, P. Gros³², J.F. Grosse-Oetringhaus³³, J.-Y. Grossiord¹¹⁶, R. Grosso³³, F. Guber⁴⁸, R. Guernane⁶⁸, B. Guerzoni²⁷, M. Guilbaud¹¹⁶, K. Gulbrandsen⁷⁷, H. Gulkanyan¹, T. Gunji¹²⁰, A. Gupta⁸⁶, R. Gupta⁸⁶, Ø. Haaland¹⁸, C. Hadjidakis⁴⁶, M. Haiduc⁵⁴, H. Hamagaki¹²⁰, G. Hamar¹²⁷, B.H. Han²⁰, L.D. Hanratty⁹⁷, A. Hansen⁷⁷, Z. Harmanová-Tóthová³⁸, J.W. Harris¹²⁸, M. Hartig⁵⁶, A. Harton¹³, D. Hasegan⁵⁴, D. Hatzifotiadiou¹⁰¹, S. Hayashi¹²⁰, A. Hayrapetyan^{33,1}, S.T. Heckel⁵⁶, M. Heide⁵⁸, H. Helstrup³⁵, A. Herghelegiu⁷⁵, G. Herrera Corral¹¹, N. Herrmann⁸⁸, B.A. Hess¹²², K.F. Hetland³⁵, B. Hicks¹²⁸, B. Hippolyte⁶², Y. Hori¹²⁰, P. Hristov³³, I. Hrivnáčová⁴⁶, M. Huang¹⁸, T.J. Humanic¹⁹, D.S. Hwang²⁰, R. Ichou⁶⁷, R. Ilkaev⁹⁴, I. Ilkiv⁷⁴,

M. Inaba¹²¹, E. Incanti²³, G.M. Innocenti²⁴, P.G. Innocenti³³, M. Ippolitov⁹⁵, M. Irfan¹⁷, C. Ivan⁹², V. Ivanov⁸¹, A. Ivanov¹²⁴, M. Ivanov⁹², O. Ivanytskyi³, A. Jacholkowski²⁶, P. M. Jacobs⁷¹, H.J. Jang⁶⁵, M.A. Janik¹²⁵, R. Janik³⁶, P.H.S.Y. Jayarathna¹¹⁷, S. Jena⁴⁴, D.M. Jha¹²⁶, R.T. Jimenez Bustamante⁵⁹, P.G. Jones⁹⁷, H. Jung⁴⁰, A. Jusko⁹⁷, A.B. Kaidalov⁵⁰, S. Kalcher³⁹, P. Kaliňák⁵¹, T. Kalliokoski⁴², A. Kalweit^{57,33}, J.H. Kang¹³⁰, V. Kaplin⁷³, A. Karasu Uysal^{33,129,66}, O. Karavichev⁴⁸, T. Karavicheva⁴⁸, E. Karpechev⁴⁸, A. Kazantsev⁹⁵, U. Kebschull⁵⁵, R. Keidel¹³¹, M.M. Khan¹⁷, P. Khan⁹⁶, K. H. Khan¹⁵, S.A. Khan¹²³, A. Khanzadeev⁸¹, Y. Kharlov⁴⁷, B. Kileng³⁵, S. Kim²⁰, M. Kim¹³⁰, M. Kim⁴⁰, J.S. Kim⁴⁰, J.H. Kim²⁰, D.W. Kim^{40,65}, B. Kim¹³⁰, D.J. Kim⁴², T. Kim¹³⁰, S. Kirsch³⁹, I. Kisel³⁹, S. Kiselev⁵⁰, A. Kisiel¹²⁵, J.L. Klay⁶, J. Klein⁸⁸, C. Klein-Bösing⁵⁸, M. Kliemant⁵⁶, A. Kluge³³, M.L. Knichel⁹², A.G. Knospe¹¹², M.K. Köhler⁹², T. Kollegger³⁹, A. Kolojvari¹²⁴, M. Kompaniets¹²⁴, V. Kondratiev¹²⁴, N. Kondratyeva⁷³, A. Konevskikh⁴⁸, R. Kour⁹⁷, V. Kovalenko¹²⁴, M. Kowalski¹¹¹, S. Kox⁶⁸, G. Koyithatta Meethalevedu⁴⁴, J. Kral⁴², I. Králik⁵¹, F. Kramer⁵⁶, A. Kravčáková³⁸, T. Krawutschke^{88,34}, M. Krelina³⁷, M. Kretz³⁹, M. Krivda^{97,51}, F. Krizek⁴², M. Krus³⁷, E. Kryshen⁸¹, M. Krzewicki⁹², Y. Kucheriaev⁹⁵, T. Kugathasan³³, C. Kuhn⁶², P.G. Kuijter⁷⁸, I. Kulakov⁵⁶, J. Kumar⁴⁴, P. Kurashvili⁷⁴, A.B. Kurepin⁴⁸, A. Kurepin⁴⁸, A. Kuryakin⁹⁴, V. Kushpil⁷⁹, S. Kushpil⁷⁹, H. Kvaerno²¹, M.J. Kweon⁸⁸, Y. Kwon¹³⁰, P. Ladrón de Guevara⁵⁹, I. Lakomov⁴⁶, R. Langoy¹⁸, S.L. La Pointe⁴⁹, C. Lara⁵⁵, A. Lardeux¹⁰⁸, P. La Rocca²⁶, R. Lea²², M. Lechman³³, G.R. Lee⁹⁷, K.S. Lee⁴⁰, S.C. Lee⁴⁰, I. Legrand³³, J. Lehnert⁵⁶, M. Lenhardt⁹², V. Lenti¹⁰⁰, H. León⁶⁰, I. León Monzón¹¹³, H. León Vargas⁵⁶, P. Lévai¹²⁷, S. Li⁷, J. Lien¹⁸, R. Lietava⁹⁷, S. Lindal²¹, V. Lindenstruth³⁹, C. Lippmann^{92,33}, M.A. Lisa¹⁹, H.M. Ljunggren³², P.I. Loenne¹⁸, V.R. Loggins¹²⁶, V. Loginov⁷³, D. Lohner⁸⁸, C. Loizides⁷¹, K.K. Loo⁴², X. Lopez⁶⁷, E. López Torres⁹, G. Løvholden²¹, X.-G. Lu⁸⁸, P. Luettig⁵⁶, M. Lunardon²⁸, J. Luo⁷, G. Luparello⁴⁹, C. Luzzi³³, K. Ma⁷, R. Ma¹²⁸, D.M. Madagadahettige-Don¹¹⁷, A. Maevskaya⁴⁸, M. Mager^{57,33}, D.P. Mahapatra⁵², A. Maire⁸⁸, M. Malaev⁸¹, I. Maldonado Cervantes⁵⁹, L. Malinina^{63,ii}, D. Mal'Kevich⁵⁰, P. Malzacher⁹², A. Mamonov⁹⁴, L. Manceau⁹⁹, L. Mangotra⁸⁶, V. Manko⁹⁵, F. Manso⁶⁷, V. Manzari¹⁰⁰, Y. Mao⁷, M. Marchisone^{67,24}, J. Mareš⁵³, G.V. Margagliotti^{22,106}, A. Margotti¹⁰¹, A. Marín⁹², C. Markert¹¹², M. Marquard⁵⁶, I. Martashvili¹¹⁹, N.A. Martin⁹², P. Martinengo³³, M.I. Martínez², A. Martínez Davalos⁶⁰, G. Martínez García¹⁰⁸, Y. Martynov³, A. Mas¹⁰⁸, S. Masciocchi⁹², M. Masera²⁴, A. Masoni¹⁰², L. Massacrier¹⁰⁸, A. Mastroserio³¹, Z.L. Matthews⁹⁷, A. Matyja^{111,108}, C. Mayer¹¹¹, J. Mazer¹¹⁹, M.A. Mazzoni¹⁰⁵, F. Meddi²⁵, A. Menchaca-Rocha⁶⁰, J. Mercado Pérez⁸⁸, M. Meres³⁶, Y. Miake¹²¹, K. Mikhailov^{63,50}, L. Milano²⁴, J. Milosevic^{21,iii}, A. Mischke⁴⁹, A.N. Mishra^{87,45}, D. Miśkowiec^{92,33}, C. Mitu⁵⁴, S. Mizuno¹²¹, J. Mlynarz¹²⁶, B. Mohanty^{123,76}, L. Molnar^{127,33,62}, L. Montaña Zetina¹¹, M. Monteno⁹⁹, E. Montes¹⁰, T. Moon¹³⁰, M. Morando²⁸, D.A. Moreira De Godoy¹¹⁴, S. Moretto²⁸, A. Morreale⁴², A. Morsch³³, V. Muccifora⁶⁹, E. Mudnic¹¹⁰, S. Muhuri¹²³, M. Mukherjee¹²³, H. Müller³³, M.G. Munhoz¹¹⁴, L. Musa³³, J. Musinsky⁵¹, A. Musso⁹⁹, B.K. Nandi⁴⁴, R. Nania¹⁰¹, E. Nappi¹⁰⁰, C. Nattrass¹¹⁹, S. Navin⁹⁷, T.K. Nayak¹²³, S. Nazarenko⁹⁴, A. Nedosekin⁵⁰, M. Nicassio^{31,92}, M. Niculescu^{54,33}, B.S. Nielsen⁷⁷, T. Niida¹²¹, S. Nikolaev⁹⁵, V. Nikolic⁹³, S. Nikulin⁹⁵, V. Nikulin⁸¹, B.S. Nilsen⁸², M.S. Nilsson²¹, F. Noferini^{101,12}, P. Nomokonov⁶³, G. Nooren⁴⁹, N. Novitzky⁴², A. Nyanin⁹⁵, A. Nyatha⁴⁴, C. Nygaard⁷⁷, J. Nystrand¹⁸, A. Ochirov¹²⁴, H. Oeschler^{57,33}, S.K. Oh⁴⁰, S. Oh¹²⁸, J. Oleniacz¹²⁵, A.C. Oliveira Da Silva¹¹⁴, C. Oppedisano⁹⁹, A. Ortiz Velasquez^{32,59}, A. Oskarsson³², P. Ostrowski¹²⁵, J. Otwinowski⁹², K. Oyama⁸⁸, K. Ozawa¹²⁰, Y. Pachmayer⁸⁸, M. Pachr³⁷, F. Padilla²⁴, P. Pagano²⁹, G. Paic⁵⁹, F. Painke³⁹, C. Pajares¹⁶, S.K. Pal¹²³, A. Palaha⁹⁷, A. Palmeri¹⁰³, V. Papikyan¹, G.S. Pappalardo¹⁰³, W.J. Park⁹², A. Passfeld⁵⁸, D.I. Patalakha⁴⁷, V. Paticchio¹⁰⁰, B. Paul⁹⁶, A. Pavlinov¹²⁶, T. Pawlak¹²⁵, T. Peitzmann⁴⁹, H. Pereira Da Costa¹⁴, E. Pereira De Oliveira Filho¹¹⁴, D. Peresunko⁹⁵, C.E. Pérez Lara⁷⁸, D. Perini³³, D. Perrino³¹, W. Peryt¹²⁵, A. Pesci¹⁰¹, V. Peskov^{33,59}, Y. Pestov⁵, V. Petráček³⁷, M. Petran³⁷, M. Petris⁷⁵, P. Petrov⁹⁷, M. Petrovici⁷⁵, C. Petta²⁶, S. Piano¹⁰⁶, A. Piccotti⁹⁹, M. Pikna³⁶, P. Pillot¹⁰⁸, O. Pinazza³³, L. Pinsky¹¹⁷, N. Pitz⁵⁶, D.B. Piyarathna¹¹⁷, M. Planinic⁹³, M. Płoskoń⁷¹, J. Pluta¹²⁵, T. Pocheptsov⁶³, S. Pochybova¹²⁷, P.L.M. Podesta-Lerma¹¹³, M.G. Poghosyan³³, K. Polák⁵³, B. Polichtchouk⁴⁷, A. Pop⁷⁵, S. Porteboeuf-Houssais⁶⁷, V. Pospíšil³⁷, B. Potukuchi⁸⁶, S.K. Prasad¹²⁶, R. Preghenella^{101,12}, F. Prino⁹⁹, C.A. Pruneau¹²⁶, I. Pshenichnov⁴⁸, G. Puddu²³, V. Punin⁹⁴, M. Putiš³⁸, J. Putschke¹²⁶, E. Quercigh³³, H. Qvigstad²¹, A. Rachevski¹⁰⁶, A. Rademakers³³, T.S. Rähä⁴², J. Rak⁴², A. Rakotozafindrabe¹⁴, L. Ramello³⁰, A. Ramírez Reyes¹¹, R. Raniwala⁸⁷, S. Raniwala⁸⁷, S.S. Räsänen⁴², B.T. Rascanu⁵⁶, D. Rathee⁸³, K.F. Read¹¹⁹, J.S. Real⁶⁸, K. Redlich^{74,61}, R.J. Reed¹²⁸, A. Rehman¹⁸, P. Reichelt⁵⁶, M. Reicher⁴⁹, R. Renfordt⁵⁶, A.R. Reolon⁶⁹, A. Reshetin⁴⁸, F. Rettig³⁹, J.-P. Revol³³, K. Reygers⁸⁸, L. Riccati⁹⁹, R.A. Ricci⁷⁰, T. Richert³², M. Richter²¹, P. Riedler³³, W. Riegler³³, F. Riggi^{26,103}, M. Rodríguez Cahuantzi², A. Rodríguez Manso⁷⁸, K. Røed^{18,21}, D. Rohr³⁹, D. Röhrich¹⁸, R. Romita^{92,107}, F. Ronchetti⁶⁹, P. Rosnet⁶⁷, S. Rossegger³³, A. Rossi^{33,28}, P. Roy⁹⁶, C. Roy⁶²,

A.J. Rubio Montero¹⁰, R. Rui²², R. Russo²⁴, E. Ryabinkin⁹⁵, A. Rybicki¹¹¹, S. Sadovsky⁴⁷, K. Šafařík³³, R. Sahoo⁴⁵, P.K. Sahu⁵², J. Saini¹²³, H. Sakaguchi⁴³, S. Sakai⁷¹, D. Sakata¹²¹, C.A. Salgado¹⁶, J. Salzwedel¹⁹, S. Sambyal⁸⁶, V. Samsonov⁸¹, X. Sanchez Castro⁶², L. Šándor⁵¹, A. Sandoval⁶⁰, M. Sano¹²¹, G. Santagati²⁶, R. Santoro^{33,12}, J. Sarkamo⁴², E. Scapparone¹⁰¹, F. Scarlassara²⁸, R.P. Scharenberg⁹⁰, C. Schiaua⁷⁵, R. Schicker⁸⁸, H.R. Schmidt¹²², C. Schmidt⁹², S. Schuchmann⁵⁶, J. Schukraft³³, T. Schuster¹²⁸, Y. Schutz^{33,108}, K. Schwarz⁹², K. Schweda⁹², G. Scioli²⁷, E. Scomparin⁹⁹, P.A. Scott⁹⁷, R. Scott¹¹⁹, G. Segato²⁸, I. Selyuzhenkov⁹², S. Senyukov⁶², J. Seo⁹¹, S. Serici²³, E. Serradilla^{10,60}, A. Sevcenco⁵⁴, A. Shabetai¹⁰⁸, G. Shabratova⁶³, R. Shahoyan³³, S. Sharma⁸⁶, N. Sharma^{83,119}, S. Rohni⁸⁶, K. Shigaki⁴³, K. Shtejer⁹, Y. Sibiriak⁹⁵, E. Sicking⁵⁸, S. Siddhanta¹⁰², T. Siemiarczuk⁷⁴, D. Silvermyr⁸⁰, C. Silvestre⁶⁸, G. Simatovic^{59,93}, G. Simonetti³³, R. Singaraju¹²³, R. Singh⁸⁶, S. Singha^{123,76}, V. Singhal¹²³, T. Sinha⁹⁶, B.C. Sinha¹²³, B. Sitar³⁶, M. Sitta³⁰, T.B. Skaali²¹, K. Skjerdal¹⁸, R. Smakal³⁷, N. Smirnov¹²⁸, R.J.M. Snellings⁴⁹, C. Sogaard^{77,32}, R. Soltz⁷², H. Son²⁰, M. Song¹³⁰, J. Song⁹¹, C. Soos³³, F. Soramel²⁸, I. Sputowska¹¹¹, M. Spyropoulou-Stassinaki⁸⁴, B.K. Srivastava⁹⁰, J. Stachel⁸⁸, I. Stan⁵⁴, G. Stefanek⁷⁴, M. Steinpreis¹⁹, E. Stenlund³², G. Steyn⁸⁵, J.H. Stiller⁸⁸, D. Stocco¹⁰⁸, M. Stolpovskiy⁴⁷, P. Strmen³⁶, A.A.P. Suaide¹¹⁴, M.A. Subieta Vásquez²⁴, T. Sugitate⁴³, C. Suire⁴⁶, R. Sultanov⁵⁰, M. Šumbera⁷⁹, T. Susa⁹³, T.J.M. Symons⁷¹, A. Szanto de Toledo¹¹⁴, I. Szarka³⁶, A. Szczepankiewicz^{111,33}, A. Szostak¹⁸, M. Szymański¹²⁵, J. Takahashi¹¹⁵, J.D. Tapia Takaki⁴⁶, A. Tarantola Peloni⁵⁶, A. Tarazona Martinez³³, A. Tauro³³, G. Tejada Muñoz², A. Telesca³³, C. Terrevoli³¹, J. Thäder⁹², D. Thomas⁴⁹, R. Tieulent¹¹⁶, A.R. Timmins¹¹⁷, D. Tlusty³⁷, A. Toia^{39,28,104}, H. Torii¹²⁰, L. Toscano⁹⁹, V. Trubnikov³, D. Truesdale¹⁹, W.H. Trzaska⁴², T. Tsuji¹²⁰, A. Tumkin⁹⁴, R. Turrisi¹⁰⁴, T.S. Tveter²¹, J. Ulery⁵⁶, K. Ullaland¹⁸, J. Ulrich^{64,55}, A. Uras¹¹⁶, J. Urbán³⁸, G.M. Urciuoli¹⁰⁵, G.L. Usai²³, M. Vajzer^{37,79}, M. Vala^{63,51}, L. Valencia Palomo⁴⁶, S. Vallero⁸⁸, P. Vande Vyvre³³, M. van Leeuwen⁴⁹, L. Vannucci⁷⁰, A. Vargas², R. Varma⁴⁴, M. Vasileiou⁸⁴, A. Vasiliev⁹⁵, V. Vechernin¹²⁴, M. Veldhoen⁴⁹, M. Venaruzzo²², E. Vercellin²⁴, S. Vergara², R. Vernet⁸, M. Verweij⁴⁹, L. Vickovic¹¹⁰, G. Viesti²⁸, Z. Vilakazi⁸⁵, O. Villalobos Baillie⁹⁷, Y. Vinogradov⁹⁴, L. Vinogradov¹²⁴, A. Vinogradov⁹⁵, T. Virgili²⁹, Y.P. Viyogi¹²³, A. Vodopyanov⁶³, K. Voloshin⁵⁰, S. Voloshin¹²⁶, G. Volpe³³, B. von Haller³³, I. Vorobyev¹²⁴, D. Vranic⁹², J. Vrláková³⁸, B. Vulpescu⁶⁷, A. Vyushin⁹⁴, V. Wagner³⁷, B. Wagner¹⁸, R. Wan⁷, D. Wang⁷, M. Wang⁷, Y. Wang⁷, Y. Wang⁸⁸, K. Watanabe¹²¹, M. Weber¹¹⁷, J.P. Wessels^{33,58}, U. Westerhoff⁵⁸, J. Wiechula¹²², J. Wikne²¹, M. Wilde⁵⁸, A. Wilk⁵⁸, G. Wilk⁷⁴, M.C.S. Williams¹⁰¹, B. Windelband⁸⁸, L. Xaplanteris Karampatsos¹¹², C.G. Yaldo¹²⁶, Y. Yamaguchi¹²⁰, H. Yang^{14,49}, S. Yang¹⁸, S. Yasnopolskiy⁹⁵, J. Yi⁹¹, Z. Yin⁷, I.-K. Yoo⁹¹, J. Yoon¹³⁰, W. Yu⁵⁶, X. Yuan⁷, I. Yushmanov⁹⁵, V. Zaccolo⁷⁷, C. Zach³⁷, C. Zampolli¹⁰¹, S. Zaporozhets⁶³, A. Zarochentsev¹²⁴, P. Závada⁵³, N. Zaviyalov⁹⁴, H. Zbroszczyk¹²⁵, P. Zelniczek⁵⁵, I.S. Zgura⁵⁴, M. Zhalov⁸¹, H. Zhang⁷, X. Zhang^{67,7}, D. Zhou⁷, Y. Zhou⁴⁹, F. Zhou⁷, J. Zhu⁷, H. Zhu⁷, J. Zhu⁷, X. Zhu⁷, A. Zichichi^{27,12}, A. Zimmermann⁸⁸, G. Zinovjev³, Y. Zoccarato¹¹⁶, M. Zynovyev³, M. Zyzak⁵⁶

Affiliation notes

ⁱ Deceased

ⁱⁱ Also at: M.V.Lomonosov Moscow State University, D.V.Skobeltzyn Institute of Nuclear Physics, Moscow, Russia

ⁱⁱⁱ Also at: University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia

Collaboration Institutes

¹ A. I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation, Yerevan, Armenia

² Benemérita Universidad Autónoma de Puebla, Puebla, Mexico

³ Bogolyubov Institute for Theoretical Physics, Kiev, Ukraine

⁴ Bose Institute, Department of Physics and Centre for Astroparticle Physics and Space Science (CAPSS), Kolkata, India

⁵ Budker Institute for Nuclear Physics, Novosibirsk, Russia

⁶ California Polytechnic State University, San Luis Obispo, California, United States

⁷ Central China Normal University, Wuhan, China

⁸ Centre de Calcul de l'IN2P3, Villeurbanne, France

⁹ Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba

¹⁰ Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

¹¹ Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City and Mérida, Mexico

- 12 Centro Fermi - Museo Storico della Fisica e Centro Studi e Ricerche “Enrico Fermi”, Rome, Italy
- 13 Chicago State University, Chicago, United States
- 14 Commissariat à l’Energie Atomique, IRFU, Saclay, France
- 15 COMSATS Institute of Information Technology (CIIT), Islamabad, Pakistan
- 16 Departamento de Física de Partículas and IGFAE, Universidad de Santiago de Compostela, Santiago de Compostela, Spain
- 17 Department of Physics Aligarh Muslim University, Aligarh, India
- 18 Department of Physics and Technology, University of Bergen, Bergen, Norway
- 19 Department of Physics, Ohio State University, Columbus, Ohio, United States
- 20 Department of Physics, Sejong University, Seoul, South Korea
- 21 Department of Physics, University of Oslo, Oslo, Norway
- 22 Dipartimento di Fisica dell’Università and Sezione INFN, Trieste, Italy
- 23 Dipartimento di Fisica dell’Università and Sezione INFN, Cagliari, Italy
- 24 Dipartimento di Fisica dell’Università and Sezione INFN, Turin, Italy
- 25 Dipartimento di Fisica dell’Università ‘La Sapienza’ and Sezione INFN, Rome, Italy
- 26 Dipartimento di Fisica e Astronomia dell’Università and Sezione INFN, Catania, Italy
- 27 Dipartimento di Fisica e Astronomia dell’Università and Sezione INFN, Bologna, Italy
- 28 Dipartimento di Fisica e Astronomia dell’Università and Sezione INFN, Padova, Italy
- 29 Dipartimento di Fisica ‘E.R. Caianiello’ dell’Università and Gruppo Collegato INFN, Salerno, Italy
- 30 Dipartimento di Scienze e Innovazione Tecnologica dell’Università del Piemonte Orientale and Gruppo Collegato INFN, Alessandria, Italy
- 31 Dipartimento Interateneo di Fisica ‘M. Merlin’ and Sezione INFN, Bari, Italy
- 32 Division of Experimental High Energy Physics, University of Lund, Lund, Sweden
- 33 European Organization for Nuclear Research (CERN), Geneva, Switzerland
- 34 Fachhochschule Köln, Köln, Germany
- 35 Faculty of Engineering, Bergen University College, Bergen, Norway
- 36 Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava, Slovakia
- 37 Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic
- 38 Faculty of Science, P.J. Šafárik University, Košice, Slovakia
- 39 Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- 40 Gangneung-Wonju National University, Gangneung, South Korea
- 41 Gauhati University, Department of Physics, Guwahati, India
- 42 Helsinki Institute of Physics (HIP) and University of Jyväskylä, Jyväskylä, Finland
- 43 Hiroshima University, Hiroshima, Japan
- 44 Indian Institute of Technology Bombay (IIT), Mumbai, India
- 45 Indian Institute of Technology Indore, Indore, India (IITI)
- 46 Institut de Physique Nucléaire d’Orsay (IPNO), Université Paris-Sud, CNRS-IN2P3, Orsay, France
- 47 Institute for High Energy Physics, Protvino, Russia
- 48 Institute for Nuclear Research, Academy of Sciences, Moscow, Russia
- 49 Nikhef, National Institute for Subatomic Physics and Institute for Subatomic Physics of Utrecht University, Utrecht, Netherlands
- 50 Institute for Theoretical and Experimental Physics, Moscow, Russia
- 51 Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovakia
- 52 Institute of Physics, Bhubaneswar, India
- 53 Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic
- 54 Institute of Space Sciences (ISS), Bucharest, Romania
- 55 Institut für Informatik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- 56 Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- 57 Institut für Kernphysik, Technische Universität Darmstadt, Darmstadt, Germany
- 58 Institut für Kernphysik, Westfälische Wilhelms-Universität Münster, Münster, Germany
- 59 Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico
- 60 Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico
- 61 Institut of Theoretical Physics, University of Wrocław
- 62 Institut Pluridisciplinaire Hubert Curien (IPHC), Université de Strasbourg, CNRS-IN2P3, Strasbourg,

- France
- ⁶³ Joint Institute for Nuclear Research (JINR), Dubna, Russia
 - ⁶⁴ Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
 - ⁶⁵ Korea Institute of Science and Technology Information, Daejeon, South Korea
 - ⁶⁶ KTO Karatay University, Konya, Turkey
 - ⁶⁷ Laboratoire de Physique Corpusculaire (LPC), Clermont Université, Université Blaise Pascal, CNRS-IN2P3, Clermont-Ferrand, France
 - ⁶⁸ Laboratoire de Physique Subatomique et de Cosmologie (LPSC), Université Joseph Fourier, CNRS-IN2P3, Institut Polytechnique de Grenoble, Grenoble, France
 - ⁶⁹ Laboratori Nazionali di Frascati, INFN, Frascati, Italy
 - ⁷⁰ Laboratori Nazionali di Legnaro, INFN, Legnaro, Italy
 - ⁷¹ Lawrence Berkeley National Laboratory, Berkeley, California, United States
 - ⁷² Lawrence Livermore National Laboratory, Livermore, California, United States
 - ⁷³ Moscow Engineering Physics Institute, Moscow, Russia
 - ⁷⁴ National Centre for Nuclear Studies, Warsaw, Poland
 - ⁷⁵ National Institute for Physics and Nuclear Engineering, Bucharest, Romania
 - ⁷⁶ National Institute of Science Education and Research, Bhubaneswar, India
 - ⁷⁷ Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
 - ⁷⁸ Nikhef, National Institute for Subatomic Physics, Amsterdam, Netherlands
 - ⁷⁹ Nuclear Physics Institute, Academy of Sciences of the Czech Republic, Řež u Prahy, Czech Republic
 - ⁸⁰ Oak Ridge National Laboratory, Oak Ridge, Tennessee, United States
 - ⁸¹ Petersburg Nuclear Physics Institute, Gatchina, Russia
 - ⁸² Physics Department, Creighton University, Omaha, Nebraska, United States
 - ⁸³ Physics Department, Panjab University, Chandigarh, India
 - ⁸⁴ Physics Department, University of Athens, Athens, Greece
 - ⁸⁵ Physics Department, University of Cape Town and iThemba LABS, National Research Foundation, Somerset West, South Africa
 - ⁸⁶ Physics Department, University of Jammu, Jammu, India
 - ⁸⁷ Physics Department, University of Rajasthan, Jaipur, India
 - ⁸⁸ Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
 - ⁸⁹ Politecnico di Torino, Turin, Italy
 - ⁹⁰ Purdue University, West Lafayette, Indiana, United States
 - ⁹¹ Pusan National University, Pusan, South Korea
 - ⁹² Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung, Darmstadt, Germany
 - ⁹³ Rudjer Bošković Institute, Zagreb, Croatia
 - ⁹⁴ Russian Federal Nuclear Center (VNIIEF), Sarov, Russia
 - ⁹⁵ Russian Research Centre Kurchatov Institute, Moscow, Russia
 - ⁹⁶ Saha Institute of Nuclear Physics, Kolkata, India
 - ⁹⁷ School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom
 - ⁹⁸ Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru
 - ⁹⁹ Sezione INFN, Turin, Italy
 - ¹⁰⁰ Sezione INFN, Bari, Italy
 - ¹⁰¹ Sezione INFN, Bologna, Italy
 - ¹⁰² Sezione INFN, Cagliari, Italy
 - ¹⁰³ Sezione INFN, Catania, Italy
 - ¹⁰⁴ Sezione INFN, Padova, Italy
 - ¹⁰⁵ Sezione INFN, Rome, Italy
 - ¹⁰⁶ Sezione INFN, Trieste, Italy
 - ¹⁰⁷ Nuclear Physics Group, STFC Daresbury Laboratory, Daresbury, United Kingdom
 - ¹⁰⁸ SUBATECH, Ecole des Mines de Nantes, Université de Nantes, CNRS-IN2P3, Nantes, France
 - ¹⁰⁹ Suranaree University of Technology, Nakhon Ratchasima, Thailand
 - ¹¹⁰ Technical University of Split FESB, Split, Croatia
 - ¹¹¹ The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland
 - ¹¹² The University of Texas at Austin, Physics Department, Austin, TX, United States
 - ¹¹³ Universidad Autónoma de Sinaloa, Culiacán, Mexico

-
- ¹¹⁴ Universidade de São Paulo (USP), São Paulo, Brazil
¹¹⁵ Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil
¹¹⁶ Université de Lyon, Université Lyon 1, CNRS/IN2P3, IPN-Lyon, Villeurbanne, France
¹¹⁷ University of Houston, Houston, Texas, United States
¹¹⁸ University of Technology and Austrian Academy of Sciences, Vienna, Austria
¹¹⁹ University of Tennessee, Knoxville, Tennessee, United States
¹²⁰ University of Tokyo, Tokyo, Japan
¹²¹ University of Tsukuba, Tsukuba, Japan
¹²² Eberhard Karls Universität Tübingen, Tübingen, Germany
¹²³ Variable Energy Cyclotron Centre, Kolkata, India
¹²⁴ V. Fock Institute for Physics, St. Petersburg State University, St. Petersburg, Russia
¹²⁵ Warsaw University of Technology, Warsaw, Poland
¹²⁶ Wayne State University, Detroit, Michigan, United States
¹²⁷ Wigner Research Centre for Physics, Hungarian Academy of Sciences, Budapest, Hungary
¹²⁸ Yale University, New Haven, Connecticut, United States
¹²⁹ Yildiz Technical University, Istanbul, Turkey
¹³⁰ Yonsei University, Seoul, South Korea
¹³¹ Zentrum für Technologietransfer und Telekommunikation (ZTT), Fachhochschule Worms, Worms, Germany